


PEROMYSCUS NEWSLETTER

Number Forty-Four



Fall 2009

Cover:

Peromyscus maniculatus bairdii blastocysts generated by harvesting zygotes 68 hours after induced ovulation and after 48 hours of culturing. Photo by G. Szalai.

Peromyscus Newsletter Number 44

Hello, All!

There are several items of particular note for this issue. First, please be aware that the USC biology department has a new email server that is not performing well. So if you send or fail to receive an email from an address with the “@biol.sc.edu” ending, please try again. We apologize for this inconvenience, but our hands are tied on this issue. In particular, this switch has also meant that the former Peromyscus@stkctr.biol.sc.edu is no longer working and may never work again, so instead please email Janet Crossland at crosslan@biol.sc.edu. Further details can be found in the *News, Comments, and Announcements* section.

Also note that the new *Peromyscus* Yahoo!Group has formed but it requires an invitation to join. I sent out invitations last year, but many people did not respond and the invitation expired. I will try sending out new invitations to subscribers in the next few months and I encourage everyone to accept the invitation to this new list-serv as it will be the fastest way for *Peromyscus* researchers to disseminate information in a timely fashion. Alternatively, you may simply email me at mammalogist@yahoo.com and I will make sure you get an invite right away. Please note that this is a private group and cannot be found on a search of Yahoo!Groups, thus you must receive and respond to the invitation in order to join.

Finally, if you have switched email addresses or are about to switch, please let me know so I can update your contact information. And as always, all suggestions to improve the Newsletter are appreciated! Everything related to *PN* should be sent to peromyscusnewsletter@yahoo.com until the server problems are fixed, hopefully in January 2010, after which the original peromyscusnewsletter@biol.sc.edu should work once again. Thanks for your understanding.

Take care and enjoy!

Julie

PEROMYSCUS NEWSLETTER is produced by the

Peromyscus Genetic Stock Center
Department of Biological Sciences
University of South Carolina
Columbia SC 29208
E-mail: crosslan@biol.sc.edu

with support, in part, from

National Science Foundation Grant # DBI-0742680

National Institutes of Health Grant # P40-RR014279

The Stock Center sponsors **PeroBase**, a comprehensive database for peromyscine rodents.

Julie L. Glenn, Editor
Peromyscus Genetic Stock Center
University of South Carolina
Columbia, SC 29208
(803) 576-5775
mammalogist@yahoo.com

Wallace D. Dawson, Editor Emeritus
Department of Biological Sciences
University of South Carolina
Columbia, SC 29208
(803) 777-3107 or (314) 835-1552
dawson@biol.sc.edu

Michael R. Felder, Director
Peromyscus Genetic Stock Center
University of South Carolina
Columbia, SC 29208
(803) 777-5135
felder@biol.sc.edu

Gabor Szalai, Associate Director
Peromyscus Genetic Stock Center
University of South Carolina
Columbia, SC 29208
(803) 576-5775
gszalai@biol.sc.edu

Michael J. Dewey, Consultant
Peromyscus Genetic Stock Center
University of South Carolina
Columbia, SC 29208
(803) 777-4132
dewey@biol.sc.edu

Janet Crossland, Staff Assistant
and Colony Manager
Peromyscus Genetic Stock Center
University of South Carolina
Columbia, SC 29208
(803) 777-3107
crosslan@biol.sc.edu

Peromyscus Genetic Stock Center Advisory Committee:

Ernest Bailey
Bruce Cushing
Courtney DeVries
Mike Hooper
Tony Schountz

Gluck Equine Research Center, U of KY
University of Akron, OH
Ohio State University
USGS, Columbia Environ Res Ctr, MO
University of Northern Colorado

Michael J. Dewey, *Ex officio*
Wallace D. Dawson, *Ex officio*

CONTENTS

<i>Peromyscus</i> Newsletter Number 44	3
News, Comments, and Announcements	6
The <i>Peromyscus</i> Genetic Stock Center	11
Schedule of User Fees	12
Stocks Available	13
Other Resources of the <i>Peromyscus</i> Stock Center	15
Spotlight on STARI: Southern Tick-Associated Rash Illness	16
Contributed Accounts	22
Population Assessment of Golden Mice in Illinois: Relative Abundance and Sympatric White-Footed Mice Denise STETSON, Jay HUBERT, and George FELDHAMER	23
Introduction of Exotic Species on Islands: <i>Peromyscus</i> Found on Shemya Island of the Far Western Aleutian Islands of Alaska Yadéeh E. SAWYER, David BANKS-RICHARDSON, S. O. MACDONALD, and Joseph A. COOK.....	27
Detecting Natural Selection in Pacific Northwest Deer Mice: An Integrative Approach Dou-Shuan YANG* and Jim KENAGY	29
Call for Data / Collaborators on Rodent Diel Activity Patterns Ivan CASTRO-ARELLANO	32
RECENT PUBLICATIONS	34

News, Comments, and Announcements

VERY IMPORTANT: USC SERVER PROBLEMS!!!

The biology department at the University of South Carolina recently switched to a new server which has been problematic. As a result, the long-held email address, Peromyscus@stkctr.biol.sc.edu no longer works and may never work again. If you need to get in contact with Stock Center personnel, please email either colony manager Janet Crossland at crosslan@biol.sc.edu or Associate Director Gabor Szalai at szalai@biol.sc.edu. The server has also not reliably sent out or delivered mail to and from the peromyscusnewsletter@biol.sc.edu account, either. Therefore, until this issue is fixed, hopefully in January, please send all Newsletter correspondence to peromyscusnewsletter@yahoo.com. Thanks so much for your understanding!



OLD ISSUES OF *PN* ON WEB SITE

All of the old issues of *Peromyscus Newsletter* are now scanned and available on the stock center web site! Thank you Janet Crossland and Shelby Carter for all those efforts!

stkctr.biol.sc.edu/Peromyscus.Newsletter.htm



RECENTLY PUBLISHED

We are thrilled to announce the long-awaited publication of the *Mammalian Species* account for *Peromyscus boylii*, the brush mouse! Thanks to Matina Kalcounis-Rueppell and Tracey Spoon for all your hard work! The species account is number 838 published in July 2009. The full citation is:

Kalcounis-Rueppell MC, and Spoon TR. 2009. *Peromyscus boylii*. *Mammalian Species*, 838, 1-14.



DR. PAUL VRANA AT USC!

We are very pleased to announce that Dr. Paul Vrana, pioneering researcher in the field of genetic imprinting and enthusiastic supporter of *Peromyscus* research, has come to the University of South Carolina! The Stock Center is thrilled to have him so close by where he can interact with us and contribute years of *Peromyscus* experience. We're so happy to have you, Paul!





OBITUARY

We are sad to report the death of Dr. Terry Yates at the age of 57. He died of brain cancer on December 11, 2007 at the University of New Mexico Health Sciences Center in Albuquerque. Dr. Yates is best known to the *Peromyscus* community for his work identifying the cause of what we now as Hantavirus Pulmonary Syndrome. Dr. Yates' collection of mice housed in the museum of Southwestern Biology was instrumental in tracking down *P. maniculatus* as the carrier of Sin Nombre Virus. So important was this discovery that The National Science Foundation named the Hanta Virus research done by Dr. Yates and his collaborator Dr. Robert Parmenter as one of NSF's Nifty 50 discoveries—projects which have had the greatest impact on the lives of Americans.

Dr. Yates served as Vice Provost for Research at the University of New Mexico in 2001, then served as Vice President for Research and Economic Development from 2004 until his recent death. He was also the Curator of Genomic Resources for the Museum of Southwestern Biology at UNM, a professor of biology and pathology, a member of the Board of Life Sciences of the National Academy of Sciences, and he helped create the Long Term Ecological Research site near Socorro. Dr. Yates was also a former external advisory committee member for the *Peromyscus* Genetic Stock Center. He contributed a great deal toward advancing our knowledge of our favorite mice and he will be greatly missed.





OBITUARY

We are also sad to report the death of Dr. Betty Horner on her birthday, April 29, 2009 at the age of 93. She received her PhD from the University of Michigan in 1948 but spent her entire academic career at Smith College. Dr. Horner interacted directly with the Stock Center on several occasions and even contributed one of our special coat color strains, golden nugget, which she described with Stock Center founder, Dr. Wally Dawson, in 1993.

In addition to her work on *Peromyscus* coat color, Dr. Horner studied topics that varied from predation on overwintering monarch butterflies to paternal care of young mice. Her interaction with the PGSC was quite close, and over her long and productive career she contributed 600 35mm slides of various *Peromyscus* and related species, many black and white pictures of *P. polionotus leucocephalus*, and detailed annotations in her many notebooks. She also donated three books to the PGSC, all of which are significant to researchers of *Peromyscus*: “Man’s Nature and Nature’s Man” and “The Biotic Provinces of North America”, both authored by Dr. Lee Dice, as well as “Report Upon Natural History Collections Made in Alaska Between the Years 1877 and 1881”. She also contributed a motion picture sequence of climbing behavior in various species and subspecies of *Peromyscus* which she made while working with Dr. Dice at Michigan, as well as a tape showing the waltzing (whirling) behavior in non-agouti *P. maniculatus*, a description of which can be found in PN28.

She was such an inspiration that Dr. Dawson featured her as our *Peromyscus* pioneer in *Peromyscus Newsletter* issue 13. Appreciation for her work, however, extended much further. In 1986 Dr. Horner was honored with a special award from the American Society of Mammalogists for her many contributions to the study of mammalogy. In addition to a certificate, she was presented with a beautifully carved wooden *Peromyscus*. She will be sorely missed.



Photograph by Dick Fish, 1986





PEROMYSCUS YAHOO! GROUP

In the last issue of PN I mentioned here the formation of a new list-serve in the form of a Google Group. Due to difficulties with that format, the new list-serve is now in the form of a Yahoo! Group, which offers many user-friendly features that I believe subscribers will find helpful, including the ability to search archives. This new group was created in an effort to improve communication within the *Peromyscus* community. For those of you who are not familiar, this is a list serve complete with searchable archives. So members may post questions to other *Peromyscus* researchers instantly. In order to maintain your privacy I have restricted the group to people who have been invited to join. Only members can post messages, read the archives, view the members list, create pages and upload files. This group and its archives does not appear in public Yahoo! search results or the directory. The only way to subscribe to this new group is by invitation only. I have tried sending invitations to everyone who also receives PN, but many email addresses are no longer valid, and the invitation expires after one month. If you would like to subscribe, please email me and I will send you an invitation! The preferred email address to use at this time is my personal email account, mammalogist@yahoo.com because the new biology server has been less than reliable and I don't want to miss anyone's message!



CLASSIC REFERENCE NOW ONLINE!

Osgood's classic reference, "Revision of the mice of the American Genus *Peromyscus*" originally published in 1909, has now been digitized by the University of South Carolina library and can be viewed at

<http://digital.tcl.sc.edu/cdm4/document.php?CISOROOT=/naf&CISOPTR=285&REC=4>

or as a pdf on the PGSC website (thanks to Paul Vrana) at

<http://stkctr.biol.sc.edu/Osgood-1909.pdf>

The full citation is:

Osgood WH. 1909. Revision of the mice of the American Genus *Peromyscus*. North American Fauna 28:1-285.






PEROMYSCUS ON NEW USC T-SHIRT!

The Department of Biological Sciences at the University of South Carolina has included a Peromyscus in its new t-shirt design!

UNIVERSITY OF SOUTH CAROLINA



ALABAMA BEACH MOUSE IN NATIONAL GEOGRAPHIC

National Geographic published a beautiful photo spread of 20 endangered and threatened species in the United States, one of which was the Alabama Beach Mouse (*P. polionotus ammobates*). To view the photo go to:

<http://ngm.nationalgeographic.com/2009/01/endangered-species/photo-map-interactive>



The 89th annual meeting of the **American Society of Mammalogists** took place June 24-28, 2009 at the University of Alaska, Fairbanks. There were an amazing 30 presentations relating to *Peromyscus*, some of which are included in the contributions section.



The 2010 annual meeting of the **American Society of Mammalogists** will take place June 11-15, 2009 at the University of Wyoming, Laramie.



USC ACCREDITATION

Congratulations to the University of South Carolina's Animal Resource Facility for once again passing inspection by the Association for Assessment and Accreditation of Laboratory Animal Care. USC has been accredited since 1984.



THE *PEROMYSCUS* GENETIC STOCK CENTER

General

The University of South Carolina has maintained a genetic stock center for *Peromyscus* (deer mice and congeneric species) since 1985. The center was established under a grant from the Living Stocks Collection Program of the National Science Foundation and continues to be supported by NSF and the NIH Biological Models and Materials Research Program. It also receives support from the University and from user fees.

The major function of the Stock Center is to provide genetically characterized types of *Peromyscus* to scientific investigators and educators. Continuation of the center is dependent upon significant external utilization, therefore potential **users are encouraged to take advantage of this resource.**

Policies and Procedures

The Stock Center maintains several categories of stocks of living animals: 1) Closed colony random-bred¹ "wild-type" stocks of seven species of *Peromyscus*. 2) Two highly inbred² stocks of "wild-type" *P. leucopus*. 3) Stocks of fifteen coat color mutations, mostly in *P. maniculatus*. 4) Stocks of eight other monogenic traits. The Stock Center operates in strict compliance with the Animal Welfare Act and is located in an AAALAC approved facility. All animal care is performed by certified technicians. Stocks are monitored regularly for presence of disease and parasites and are free of hantavirus and 15 murine viruses.

The Stock Center also provides blood, organs, tissues, fetuses, skins and other biological materials from *Peromyscus*. The Stock Center operates a Molecular Bank where selected genomic libraries and probes are available. Other resources include a reference collection of more than 2,500 reprints of articles on peromyscine rodents, copies of which may be provided. The Stock Center is the primary sponsor of **PeroBase**, an on-line database dedicated to information regarding *Peromyscus* and closely related species.

Sufficient animals of the mutant types generally can be provided to initiate a breeding stock. Somewhat larger numbers, up to about 50 animals, can be provided from the wild-type stocks. Animals requested in greater numbers frequently require a "breed-up" charge and some delay in shipment.

Orders and Pricing

A user fee is charged for animals or materials provided by the Stock Center. A schedule of fees is shown on the next page. Fees vary with species and type of service provided. User assumes the cost of all shipment. Animals lost in transit are replaced without charge. Tissues, blood, skins, etc. are supplied at a modest fee that includes technician time. Arrangements for special orders will be negotiated. Billing will be submitted upon satisfactory delivery. **Write or call for details or special requirements.**

SCHEDULE OF USER FEES

Item	Academic and Government	Commercial
MATURE ANIMALS (each)		
Wild-type Stocks		
Smaller species (<i>P. maniculatus</i> , <i>P. polionotus</i> , <i>P. leucopus</i> , <i>P. eremicus</i>)	\$ 25.00	\$40.00
Larger species (<i>P. californicus</i> , <i>P. melanophrys</i> , <i>P. aztecus</i>)	33.00	45.00
Mutant and Inbred Stocks	33.00	45.00
Pregnant females (Smaller species)	45.00	55.00
(Larger species)	60.00	70.00
Special Attention (Diet, <i>etc.</i>)	45.00	55.00
F ₁ Species Hybrids	35.00	45.00
TISSUE SAMPLES (Per sample) Cost of the animal + lab fee + materials		
Solid		
Fluid (Blood, urine, saliva, <i>etc.</i>) per ml		
Flat skins (each)		
MOLECULAR MATERIALS		
Extracted DNA, 10 µg	20.00	
PCR Primers (500 µl @ 10 µM)	10.00	
Genomic & cDNA libraries	300.00	

OTHER CHARGES

Shipping costs = actual shipper's charges plus cost of mouse containers, packaging.

Lab fee for sample preparation (\$35.00/hr).

Breed-up fees (for orders exceeding 50 animals) = *per diem* cage charges X cages required.

STOCKS AVAILABLE

WILD TYPE STOCKS

ORIGIN

<i>P. maniculatus bairdii</i> (BW Stock) Deer Mouse	Closed colony bred in captivity since 1948. Descended from 40 ancestors wild-caught near Ann Arbor MI.
<i>P. maniculatus sonoriensis</i> (SM2 Stock) Sonoran Deer Mouse	Derived from about 50 animals wild-caught by Jack Hayes in 1995 near White Mountain Research Station CA.
<i>P. polionotus subgriseus</i> (PO Stock) Oldfield Mouse	Closed colony since 1952. Derived from 21 ancestors wild-caught in Ocala Nat'l. Forest FL. High inbreeding coefficient.
<i>P. leucopus</i> (LL Stock) White-footed Mouse	Derived from 38 wild ancestors captured between 1982 and 1985 near Linville NC.
<i>P. californicus insignis</i> (IS Stock) California Mouse	Derived from about 60 ancestors collected between 1979 and 1987 in Santa Monica Mts. CA.
<i>P. aztecus</i> (AM Stock) Aztec Mouse	Derived from animals collected on Sierra Chincua Michoacan, Mexico in 1986.
<i>P. melanophrys</i> (XZ Stock) Plateau Mouse	Derived from animals collected between 1970 and 1978 from Zacatecas, Mexico and bred by R. Hill.
<i>P. eremicus</i> (EP Stock) Cactus Mouse	Originated from 10-12 animals collected at Tucson AZ in 1993.

INTERSPECIFIC HYBRIDS

<i>P. maniculatus</i> X <i>P. polionotus</i> F ₁ Hybrids	Bred by special order.
<i>P. leucopus</i> X <i>P. gossypinus</i> F ₁ Hybrids	Sometimes available by special arrangement.

³COAT COLORS

ORIGINAL SOURCE

Blonde <i>bln/bln</i>	Mich. State U. colony (Pratt and Robbins, 1982)
Albino <i>c/c</i>	Sumner's albino deer mice (Sumner, 1922)
Ashy <i>ahy/ahy</i>	Wild-caught in Oregon ~ 1960 (Teed et al., 1990)
⁴ Brown <i>b/b</i>	Huestis stocks (Huestis and Barto, 1934)
California blonde <i>cfb/cfb</i>	Santa Cruz I., Calif., stock (Roth and Dawson, 1996)
Dominant spotting <i>S/+</i>	Wild caught in Illinois (Feldman, 1936)
Golden nugget <i>b^{gn}/b^{gn}</i>	Wild caught <i>P. leucopus</i> (Horner and Dawson, 1993)
Ivory <i>i/i</i>	Wild caught in Oregon (Huestis, 1938)
Platinum <i>plt/plt</i>	Barto stock at U. Mich. (Dodson et al., 1987)
⁴ Silver <i>sil/sil</i>	Huestis stock (Huestis and Barto, 1934)
Tan streak <i>tns/tns</i>	Clemson U. stock from NC (Wang et al., 1993)
Variable white <i>Vw/+</i>	Mich. State U. colony (Cowling et al., 1994)
Wide-band agouti <i>A^{Nb}/a</i>	Natural polymorphism U. Mich. (McIntosh, 1954)

OTHER MUTATIONS AND VARIANTS

Alcohol dehydrogenase negative	<i>Adh⁰/Adh⁰</i>	South Carolina BW stock (Felder, 1975)
Alcohol dehydrogenase positive	<i>Adh^f/Adh^f</i>	South Carolina BW stock (Felder, 1975)
Boggler	<i>bgl/bgl</i>	Blair's <i>P. m. blandus</i> stock (Barto, 1955)
Cataract-webbed	<i>cwb/cwb</i>	From Huestis stocks (Anderson and Burns, 1979)
Epilepsy	<i>epl/epl</i>	U. Michigan <i>P. m. artemisiae</i> stock (Dice, 1935)
Hairless-1	<i>hr-1/hr-1</i>	Sumner's hairless mutant (Sumner, 1924)
Hairless-2	<i>hr-2/hr-2</i>	Egoscue's hairless mutant (Egoscue, 1962)
Juvenile ataxia	<i>ja/ja</i>	U. Michigan stock (Van Ooteghem, 1983)
Enzyme variants		Wild type stocks provide a reservoir of variants (Dawson, 1983)

¹ "Random bred" without deliberate selection, sib-sib matings avoided. ² Inbred lines bred by sib-sib and/or parent-offspring mating for 21 generations or more. ³ Unless otherwise noted, mutations are in *P. maniculatus*. ⁴ Available only as silver/brown double recessive.

Other Resources of the *Peromyscus* Stock Center

Highly inbred *P. leucopus* (I₃₀₊) are available as live animals or as frozen tissues.
Two lines developed by George Smith (UCLA) are currently maintained by the Stock Center.

Limited numbers of other stocks are on hand, but not currently available. Inquire.

Preserved or frozen specimens of types given in the above tables.

Flat skins of mutant or wild-type coat colors of any of the stocks listed above.

Reference library of more than 2500 reprints of research papers, articles and reports on *Peromyscus*. Single copies of individual articles can be photocopied and mailed. Please limit requests to not more than five articles at any given time. There will be a charge of 10 cents per photocopied page after the initial 20 pages.

Photocopies of back issues of *Peromyscus Newsletter* (\$5 ea.) or single original back copies, when still available, without charge.

Materials are available through the *Peromyscus* Molecular Bank of the Stock Center. Allow two weeks for delivery. Included is purified DNA or frozen tissues of any of the stocks listed above. Several genomic libraries and a variety of molecular probes are available. (Inquire for more information)

For additional information or details about any of these mutants, stocks or other materials contact: Janet Crossland, Colony Manager, Peromyscus Stock Center, (803) 777-3107, e-mail crosslan@biol.sc.edu

PLEASE CALL WITH INQUIRIES

***Peromyscus* Genetic Stock Center
University of South Carolina
Columbia, SC 29208
(803) 777-3107
(803) 777-1212
FAX (803) 576-5780
crosslan@biol.sc.edu
<http://stkctr.biol.sc.edu>**

Spotlight on STARI: Southern Tick-Associated Rash Illness

Although Lyme disease is not generally considered common in the southeastern United States, another disease with similar symptoms is gaining notoriety. First recognized in the late 1980s, the disease is called Southern Tick-Associated Rash Illness, or STARI for short. Sometimes it is referred to as Masters disease after Dr. Edwin Masters, a Missouri physician who brought the unusual rashes he saw to the attention of the Centers for Disease Control and led to the first descriptions of STARI. Below is a bit of information about this newly recognized disease which, like Lyme disease, may infect *Peromyscus*.

Etiologic Agent

Unknown, but thought to be a spirochete like *Borrelia burgdorferi*, the spirochete causing Lyme disease. *B. burgdorferi* is likely not the cause as it has never been cultured from patients showing STARI symptoms, even though it is easily cultured from Lyme patients. However, antibodies to *B. burgdorferi* proteins can be elicited when *B. burgdorferi* extracts are mixed with serum from STARI patients. Thus, the etiologic agent is believed to be very similar to *B. burgdorferi*. One candidate has been *B. lonestari* which has been detected by polymerase chain reaction in the skin lesion of one patient, however, a larger study was unsuccessful in detecting *B. lonestari* in other patients. Thus, the etiologic agent has remained elusive.

Transmission Vector


Transmission to humans is generally believed to occur via a bite from a Lone Star tick, *Amblyomma americanum*, one of the most common ticks known to feed on humans. This contrasts with Lyme disease which is transmitted from the bite of *Ixodes* ticks (e.g., *Ixodes scapularis*, the deer tick, and *Ixodes pacificus*, the western black-legged tick).



Lone Star tick nymph and adult female

Distribution


As the range of the Lone Star tick expands northward, so is the presence of STARI. In the northeastern US where Lyme disease is common, distinguishing STARI from Lyme disease has proven difficult.



Distribution of the Lone Star tick (from Masters et al., 2008)

Symptoms

Within seven days of being bitten by a Lone Star tick, patients note an expanding "bull's-eye" skin rash that is similar in appearance to the erythema migrans of Lyme disease (note, however, that not all Lyme disease rashes present as a bull's-eye pattern). Compared to Lyme disease, STARI rashes are generally smaller and less variable in shape with a clearer center, making them appear much more like a bull's-eye than typical Lyme rashes. Patients may also complain of fever, headache, stiff neck, myalgias, and joint pain, however, these complaints are usually less frequent and less severe than in patients diagnosed with Lyme disease. Similarly, although some physicians have noted post-rash neurologic and cardiac problems in STARI patients, these cases are again much rarer than reported for Lyme disease. Because Lone Star ticks are also known to be vectors of Q fever and human monocytic ehrlichiosis there is always the possibility of coinfection, which has already been documented in the scientific literature.



Typical bulls-eye rashes from STARI patients (from Masters et al., 2008)

Diagnosis

Because the etiologic agent of STARI has yet to be cultured from the rash and positively identified, diagnosis is based solely on symptoms, particularly the presence of an expanding circular rash, and ideally a history of a known bite from a Lone Star tick.

Treatment

Currently, patients diagnosed with STARI are treated with antibiotics in a similar fashion as if they had Lyme disease. Doxycycline or amoxicillin administered for 2-3 weeks is commonly prescribed and most patients respond well if treated early.

Relevant Literature

- Barbour AG, Maupin GO, Teltow GJ, Carter CJ, Piesman J. 1996. Identification of an uncultivable *Borrelia* species in the hard tick *Amblyomma americanum*: possible agent of a Lyme disease-like illness. *Journal of Infectious Diseases* 173:403-409.
- Blanton L, Keith B, Brzezinski W. 2008. Southern tick-associated rash illness: Erythema migrans is not always Lyme disease. *Southern Medical Journal* 101:759-760.
- Burkot TR, Mullen GR, Anderson R, Schneider BS, Happ CM, Zeidner NS. 2001. *Borrelia lonestari* DNA in adult *Amblyomma americanum* ticks, Alabama. *Emerging Infectious Diseases* 7:471-473.
- Campbell GL, Paul WS, Schriefer ME, Craven RB, Robbins KE, Dennis DT. 1995. Epidemiologic and diagnostic studies of patients with suspected early Lyme disease, Missouri, 1990-1993. *Journal of Infectious Diseases* 172:470-480.
- Felz MW, Chandler FW Jr, Oliver JH Jr, Rahn DW, Schriefer ME. 1999. Solitary erythema migrans in Georgia and South Carolina. *Archives of Dermatology* 135:1317-1326.
- Fukunaga M, Okada K, Nakao M, Konishi T, Sato Y. 1996. Phylogenetic analysis of *Borrelia* species based on flagellin gene sequences and its application for molecular typing of Lyme disease borreliae. *International Journal of Systematic Bacteriology* 46:898-905.
- Georgia Department of Human Resources, Division of Public Health, Epidemiology Branch. 2001. Tick Bites and Erythema Migrans in Georgia: It Might NOT be Lyme Disease! *Georgia Epidemiology Report* 2001 Aug,17:1-3.
- Haddad FA, Schwartz I, Liveris D, Wormser GP. 2005. A skin lesion in a patient from Kentucky-Southern tick-associated rash illness (STARI). *Clinical Infectious Diseases* 40:475-476.
- Hair JA, Bowman JL. 1986. Behavioral ecology of *Amblyomma americanum* (L.). Pp. 407 in *Morphology, physiology, and behavioral biology of ticks* (Sauer RJ and Hair HA, eds). West Sussex England: Ellis Horwood Limited, 1986:407.
- James AM, Liveris D, Wormser GP, Schwartz I, Montecalvo MA, Johnson BJB. 2001. *Borrelia lonestari* infection after a bite by an *Amblyomma americanum* tick. *Journal of Infectious Diseases* 183:1810-4.

- Kirkland KB, Klimko TB, Meriwether RA, Schriefer M, Levin M, Levine J, MacKenzie WR, Dennis DT. 1997. Erythema migrans-like rash illness at a camp in North Carolina: a new tick-borne disease? *Archives of Internal Medicine* 157:2635-2641.
- Luckhart S, Mullen GR, Durden LA, Wright JC. 1992. *Borrelia* species in ticks recovered from white-tailed deer in Alabama. *Journal of Wildlife Diseases* 28:449-452.
- Masters E, Granter S, Duray P, Cordes P. 1998. Physician-diagnosed erythema migrans and erythema migrans-like rashes following Lone Star tick bites. *Archives of Dermatology* 134:955-960.
- Masters EJ, Grigery CN, Masters RW. 2008. STARI, or Masters Disease: Lone Star Tick–Vectored Lyme-like Illness. *Infectious Disease Clinics of North America* 22:361-376.
- Melski JW. 1999. Language, logic, and Lyme disease. *Archives of Dermatology* 135:1398-1400.
- Moody EK, Barker RW, White JL, Crutcher JM. 1998. Ticks and tick-borne diseases in Oklahoma. *Journal of the Oklahoma State Medical Association* 91:438-445.
- Oliver JH Jr, Kollars TM Jr, Chandler FW Jr, James AM, Masters EJ, Lane RS, Huey LO. 1998. First isolation and cultivation of *Borrelia burgdorferi* sensu lato from Missouri. *Journal of Clinical Microbiology* 36:1-5.
- Philipp MT, Masters E, Wormser GP, Hogrefe W, Martin D. 2006. Serologic evaluation of patients from Missouri with erythema migrans-like skin lesions with the C-6 Lyme test. *Clinical and Vaccine Immunology* 13:1170-1171.
- Piesman J, Happ CM. 1997. Ability of the Lyme disease spirochete *Borrelia burgdorferi* to infect rodents and three species of human-biting ticks (blacklegged tick, American dog tick, lone star tick) Acari:Ixodidae). *Journal of Medical Entomology* 34:451-456.
- Piesman J, Sinsky RJ. 1988. Ability of *Ixodes scapularis*, *Dermacentor variabilis*, and *Amblyomma americanum* (Acari: Ixodidae) to acquire, maintain, and transmit Lyme disease spirochetes (*Borrelia burgdorferi*). *Journal of Medical Entomology* 25:336-339.
- Ras NM, Lascola B, Postic D, Cutler SJ, Rodhain F, Baranton G, Raoult D. 1996. Phylogenesis of relapsing fever *Borrelia* spp. *International Journal of Systematic Bacteriology* 46:859-865.

- Rich SM, Armstrong PM, Smith RD, Telford SR 3rd. 2001. Lone star tick-infecting borreliae are most closely related to the agent of bovine borreliosis. *Journal of Clinical Microbiology* 39:494-497.
- Ryder JW, Pinger RR, Glancy T. 1992. Inability of *Ixodes cookei* and *Amblyomma americanum* nymphs (Acari: Ixodidae) to transmit *Borrelia burgdorferi*. *Journal of Medical Entomology* 29:525-530.
- Stromdahl EY, Evans SR, O'Brien JJ, Gutierrez AG. 2001. Prevalence of infection in ticks submitted to the human tick test kit program of the U.S. Army Center for Health Promotion and Preventive Medicine. *Journal of Medical Entomology* 38:67-74.
- Taft SC, Miller MK, Wright SM. 2005. Distribution of Borreliae among ticks collected from eastern states. *Vector-Borne and Zoonotic Diseases* 5:383-389.
- Varela AS, Luttrell MP, Howerth EW, Moore VA, Davidson WR, Stallknecht DE, Little SE. 2004. First culture isolation of *Borrelia lonestari*, putative agent of southern tick-associated rash illness. *Journal of Clinical Microbiology* 42:1163-1169.
- Wormser GP, Masters E, Liveris D, Nowakowski J, Nadelman RB, Holgren D, Bittker A, Cooper D, Wang G, Schwartz I. 2005. Microbiologic evaluation of patients from Missouri with erythema migrans. *Clinical Infectious Diseases* 40:423-8.
- Wormser GP, Masters E, Nowakowski J, McKenna D, Holgren D, Ma K, Ihde L, Cavaliere LF, Nadelman RB. 2005. Prospective clinical evaluation of patients from Missouri and New York with erythema migrans-like skin lesions. *Clinical Infectious Diseases* 41:958-65.





NOTICE

PEROMYSCUS NEWSLETTER IS NOT A FORMAL SCIENTIFIC PUBLICATION.

THEREFORE...

INFORMATION AND DATA IN THE CONTRIBUTIONS SECTION SHOULD NOT BE CITED OR USED WITHOUT PERMISSION OF THE CONTRIBUTOR.

THANK YOU!



Population Assessment of Golden Mice in Illinois: Relative Abundance and Sympatric White-Footed Mice

Denise STETSON*, Jay HUBERT, and George FELDHAMER

Department of Zoology, Southern Illinois University, Carbondale, IL 62901-6501

*Corresponding author: denise.stetson@gmail.com

Abstract

Golden mice (*Ochrotomys nuttalli*) are on the periphery of their range in Illinois where they are a state-listed species because of limited distribution and low population density. This project was initiated to assess relative abundance and distribution of *O. nuttalli* in Illinois in comparison with populations throughout the core of their geographic range. We live trapped small mammals at 24 sites in southern Illinois; each site (7 x 7 grids) was sampled twice for 5-day periods from April-October 2008. Golden mice were trapped on 21 of the sites (mean = 4.13 individuals; range 1-15). White-footed mice (*Peromyscus leucopus*) occurred on all sites (mean = 10.04; range 2-21). We expected an inverse relationship in abundance between the two species because of interspecific competition. However, there was a significant positive relationship considering new animals for each sample period ($F = 16.32$, $df = 46$, $P < 0.0002$) as well as minimum number known alive ($F = 8.69$, $df = 46$, $P < 0.005$). Positive relationships may have resulted from sampling bias that favored golden mice because traps were placed above ground as often as possible. From May through October 2009, 24 sites throughout the southeast U.S will be sampled for comparison with Illinois populations.

Introduction

Golden mice are often locally common in the southeastern United States—the core of their geographic range—where they are considered “secure.” However, in states on the periphery of their range, including Illinois, Missouri, Texas and West Virginia, they are state listed as threatened or an equivalent (Feldhamer and Morzillo 2008). Listings of golden mice may be a result of their actual rarity, or just perceived rarity due to highly elusive behavior, spatial clustering of populations or temporal shifts in overall abundance. The purpose of this project is to determine if the listing of *O. nuttalli* in Illinois is warranted. Populations of golden mice were quantitatively assessed throughout the southern part of Illinois to determine their relative abundance and site occupancy. Results will be compared to populations from the core of the range in the southeast. If results are similar, *O. nuttalli* will be proposed for removal from the threatened list in Illinois. The initial data from trapping in Illinois offered an opportunity to examine relative abundance between golden mice and white-footed mice (*Peromyscus*

leucopus). These sympatric species exhibit a high degree of niche overlap, and are considered by many investigators to be competitors in shared habitats.

Methods

Twenty-four study sites throughout Illinois were selected using GAP analysis and GIS data layers to identify ideal habitat for golden mice. All twenty-four sites were randomly selected on public land (Shawnee National Forest, Crab Orchard National Wildlife Refuge, and Cypress Creek National Wildlife Refuge) and within 0.4 km of a road for easy accessibility.

Live trapping occurred from spring 2008 through early autumn 2008. Traps were set concurrently on 3 grids at a time for 5 consecutive nights—49 Sherman traps in a 70m x 70m grid at each site. Traps were placed in elevated locations when possible in the best available golden mouse habitat. From 30 to 49 traps were elevated on each of the 24 sites. Of 1176 traps (49 per site x 24 sites), 926 were elevated (76.7%). Traps contained fiberfill batting and were baited with sunflower seeds and cracked corn. Captured animals were marked with Monel ear tag and 1.5 mm ear tissue biopsy removed for later genetic analyses. We recorded species, gender, weight, reproductive status, station, and trap location (elevated or ground). Data were analyzed using JMP IN 4.0.2.

Results

Total captures: golden mice = 99 individuals caught 190 times (174 times in elevated traps = 91.6%). White-footed mice = 244 individuals caught 565 times (465 in elevated traps = 82.3%)

Accounting for most traps being elevated, golden mice were caught significantly more often in elevated traps ($\chi^2 = 23.53$, $df = 1$, $P < 0.00001$), as were white-footed mice ($\chi^2 = 9.91$, $df = 1$, $P < 0.005$).

For all 24 sites (pooled data): regression of golden mice caught in elevated traps vs. number of elevated traps on site was significant ($F = 5.02$, $df = 22$, $P < 0.036$). This relationship was still significant when accounting for white-footed mice caught in ground traps [i.e., did more w-footed mice on ground force golden mice to elevated traps? No] ($F = 4.29$, $df = 21$, $P < 0.027$). The number of golden mice caught in elevated traps was not related to the number of w-footed mice caught in ground traps ($F = 0.18$, $df = 22$, $P < 0.676$).

There was no relationship between w-footed mice caught in elevated traps and the number of elevated traps ($F = 0.28$, $df = 22$, $P < 0.600$). There was a significant relationship between number of w-footed mice caught on the ground and the number of ground traps ($F = 5.84$, $df = 22$, $P < 0.024$). The number of golden mice caught in ground traps was not related to the number of traps on the ground ($F = 1.47$, $df = 22$, $P < 0.238$). There was still no relationship when

corrected for number of white-footed mice captured in ground traps at same time ($F = 0.84$, $df = 21$, $P < 0.444$).

There was a significant POSITIVE relationship between golden mice and white-footed mice captured in elevated traps during same period (pooled data) ($F = 22.07$, $df = 22$, $P < 0.0001$), and for all sites on a daily basis ($F = 91.22$, $df = 238$, $P < 0.0001$).

Discussion

Over twice as many individual white-footed mice were caught almost 3X more often as golden mice is consistent with the idea that *O. nuttalli* is generally uncommon with population densities lower than sympatric species of *Peromyscus* (Feldhamer and Morzillo 2008; Rose 2008). Nonetheless, the number of golden mice we caught was greater than expected on the periphery of their range. Golden mice are arboreal, but even accounting for the elevated trap bias we caught them even more often than expected above ground. Arboreal captures of golden mice were not affected on our study sites by the number of white-footed mice captured on the ground. We expected an inverse relationship between golden mice and white-footed mice due to interspecific competition. Based on the number of arboreal captures, however, there was no apparent interspecific competitive interaction between these species on our sites. Both species were trapped significantly more often in elevated traps and showed a positive relationship in total captures. It appears that *O. nuttalli* and *P. leucopus* in southern Illinois are both responding to better arboreal habitat versus ground habitat. We can contrast our results with those of Feldhamer and Maycroft (1992) and others who found significant inverse relationships in abundance between these species (see Christopher and Cameron 2008). The role of competition between golden mice and white-footed mice certainly reflects regional and site-specific microhabitat impacts, as well as a mosaic of other interacting factors—as such, generalizations and firm conclusions remain problematic.

References

- Christopher CC, Cameron GN. 2008. Community ecology of the golden mouse. Pp 59-79 in *The golden mouse: ecology and conservation* (GW Barrett and GA Feldhamer, eds). Springer, New York.
- Feldhamer GA, Maycroft KA. 1992. Unequal capture response of sympatric golden mice and white-footed mice. *American Midland Naturalist* 128:407-410.
- Feldhamer GA, Morzillo AT. 2008. Relative abundance and conservation: is the golden mouse a rare species? Pp 117-133 in *The golden mouse: ecology and conservation* (GW Barrett and GA Feldhamer, eds). Springer, New York.

Rose RK. 2008. Population ecology of the golden mouse. Pp 39-58 in The golden mouse: ecology and conservation (GW Barrett and GA Feldhamer, eds). Springer, New York.



Introduction of Exotic Species on Islands: *Peromyscus* Found on Shemya Island of the Far Western Aleutian Islands of Alaska

Yadéeh E. SAWYER*, David BANKS-RICHARDSON, S. O. MACDONALD, and Joseph A. COOK

Department of Biology and Museum of Southwestern Biology, University of New Mexico, Albuquerque, NM 87131 USA.

*Corresponding Author: yadeeh@unm.edu


Shemya Island (Fig. 1, red triangle), one of the Semichi Islands of the Near Aleutian Islands group, now harbors an expanding population of deermice, genus *Peromyscus* (MacDonald and Cook 2009). This population was first discovered and a sample taken in 1978 by D. D. Gibson, University of Alaska Museum of the North, during the course of his field work there. Given the monumental conservation issue of island exotics in general, their presence on this remote North Pacific island, far from the known range limits of the genus, has generated questions surrounding their origin and identity.

To begin answering these questions, we collected mitochondrial cytochrome *b* sequence data from 2 individuals from Shemya. For comparison, we obtained 47 additional sequences of *Peromyscus* from GenBank.

Preliminary analysis suggests a closer genetic relationship with the Rocky Mountain States lineage (Dragoo et al., 2006) of putative *P. maniculatus* in the western Lower 48 states (Fig. 1, orange hatched range). The geographically proximate *P. maniculatus* population occurs in Atlin, British Columbia (Fig. 1, Blue circle), however, they, and all other individuals of both *P. maniculatus* and *P. keeni* from elsewhere in Alaska and central and upper British Columbia, appear to be more genetically distant. These findings suggest an anthropogenic introduction of *P. maniculatus* to this island.

The discovery of the introduced *P. maniculatus* population on Shemya Island reinforces the need for more effective plans to prevent undesired transport of exotic animals, particularly to island localities (Ebbert and Byrd 2002).

We plan to collect *cytb* and nuclear introns sequences from additional individual mice from the Shemya Island population captured over a span of 15 years to further test their origin and assess whether they are derived from a single introduction event. Knowing the origin and number of introductions of the Shemya mice will provide information regarding invasions of this species in the Aleutian Islands and elsewhere.



References

- Ebbert SE, Byrd GV. 2002. Eradications of invasive species to restore natural biological diversity on Alaska Maritime National Wildlife Refuge. Pp. 102-109 in Turning the tide: the eradication of invasive species (CR Veitch and MN Clout, eds). IUCN SSC Invasive Species Specialist Group. IUCN, Gland, Switzerland and Cambridge, UK.
- MacDonald SO, Cook JA. 2009. Recent Mammals of Alaska. University of Alaska Press, Fairbanks.
- Dragoo J, Lackey A, Moore K, Lessa E, Cook J, Yates T. 2006. Phylogeography of the deer mouse (*Peromyscus maniculatus*) provides a predictive framework for research on hantaviruses. Journal of General Virology 87:1997-2003.



Detecting Natural Selection in Pacific Northwest Deer Mice: An Integrative Approach

Dou-Shuan YANG* and Jim KENAGY

Burke Museum and Department of Biology, University of Washington, Seattle, WA, 98195

*Corresponding author: dsyang@u.washington.edu

We investigated a previously identified mtDNA genetic break in deer mice, *Peromyscus maniculatus* (Awise 1979a; Lansman *et al.* 1983; Dragoo *et al.* 2006) using both mtDNA sequences and nuclear genomic markers. We analyzed spatial variation in a 491-base-pair fragment of the mtDNA control region from 455 samples distributed across a north-south transect of 2,000 km in Western North America (Figure 1). We localized the extent of the mtDNA break to a 300 km region in southern Oregon (Figure 2). To determine whether the mtDNA break was reflected in the nuclear genome and thus represented reproductive isolation, we then compared spatial variation in 13 nuclear microsatellites of 95 individuals surrounding the mtDNA break. Using a canonical correlation analysis we found that nuclear genomic variation was not correlated with mtDNA differentiation (Figure 3), and thus did not represent reproductive isolation. The contrasting patterns of variation in mtDNA and nuclear DNA are consistent with a hypothesis of historic genetic drift that occurred in isolated Pleistocene refugia combined with recent gene flow between these formerly isolated refugial populations. A Mantel test of genetic vs. geographic distance revealed that recent gene flow between deer mouse populations has been high. Thus, we conclude that past vicariant events associated with Pleistocene climate changes together with recent gene flow have created the observed intra-specific cytonuclear discordance in Western North American *P. maniculatus*. For further detail, please see our article in press in *Molecular Ecology*.

References

- Awise JC, Lansman RA, Shade RO (1979a) The use of restriction endonucleases to measure mitochondrial DNA sequence relatedness in natural populations. I. Population structure and evolution in the genus *Peromyscus*. *Genetics*, 92, 279-295.
- Lansman RA, Awise JC, Aquadro CF, Shapira JF, Daniel SW (1983) Extensive genetic variation in mitochondrial DNA's among geographic populations of the deer mouse, *Peromyscus maniculatus*. *Evolution*, 37, 1-16.
- Dragoo JW, Lackey JA, Moore KE *et al.* (2006) Phylogeography of the deer mouse (*Peromyscus maniculatus*) provides a predictive framework for research on hantaviruses. *Journal of General Virology*, 87, 1997-2003.




Figure 1




Figure 2




Figure 3



Call for Data / Collaborators on Rodent Diel Activity Patterns

Ivan CASTRO-ARELLANO*

*Corresponding Author: ivan.castro@uconn.edu

In recent times the ecological significance of time as a mediator of interactions has been recognized as important (Kronfeld-Schor & Dayan 2003). To address if an assemblage has more or less overlap in their activities as compared to random expectation I developed a simulation model for time as a resource (Castro-Arellano et al, submitted). Time occurs as an ordered resource that emerges as cyclical phenomena (e.g., diel phases, yearly seasons, etc.) and as such needs a specific randomization procedure. This model has been successfully applied to a series of bat (Castro-Arellano et al 2009; Presley et al 2009a; Presley et al 2009b) and rodent assemblages (Castro-Arellano and Lacher 2009). In a diverse rodent assemblage examined, species show less overlap than expected by random processes, thus pointing to the possible role of time as a means for species segregation. To inquire about the generality of this pattern I am seeking collaborators for the analysis of temporal activity patterns of non-volant small mammals (e.g., rodents, marsupials, insectivores). Historically, few researchers have collected the necessary data to evaluate temporal activity for small mammals, however, with the advent of new technologies (thermal video recording, PIT tagging, video monitoring, etc.) the activity patterns of animals may be recorded while collecting data for other purposes. I am seeking activity data from small mammal assemblages that have 3 or more species from anywhere in the world. All contributors of data would be co-authors on the resulting manuscripts. If enough data is gathered it may be possible to address several questions, including:

- 1) What is the prevalence of temporal niche partitioning among coexisting species of small mammals?
- 2) What is the relationship between species richness and degree of temporal niche partitioning among species of small mammals?
- 3) Is temporal activity a species characteristic or do species change their times of activity based on species composition of local assemblages (i.e. are measured activity patterns representative of fundamental or realized niche space)?

Data could have been collected by different means but as long as it can be assigned to 1 or 2 hr time blocks it would be useful for this type of analysis. Larger sample sizes are better able to characterize activity patterns, therefore, activity patterns based on at least 30 individuals are more reliable and more confidently included in analyses. For more information about the simulation

model you can contact me directly (ivan.castro@uconn.edu) or visit the following website:

<http://hydrodictyon.eeb.uconn.edu/people/willig/Research/activity%20pattern.html>

References

Castro-Arellano I, Lacher Jr TE, Willig MR, Rangel TF. (submitted) Assessment of assemblage-wide temporal-niche segregation using null models.

Castro-Arellano I, Lacher Jr TE. (in press) Temporal niche segregation in two rodent assemblages of subtropical Mexico. *Journal of Tropical Ecology*.

Castro-Arellano I, Presley SJ, Willig MR, Wunderle JM, Saldanha LN. 2009. Reduced-impact logging and temporal activity of understory bats in lowland Amazonia. *Biological Conservation* 142:2131-2139.

Kronfeld-Schor N, Dayan T. 2003. Partitioning of time as an ecological resource. *Annual Review of Ecology, Evolution and Systematics* 34:153-181.

Presley SJ, Willig MR, Castro-Arellano I, Weaver SC. 2009a. Effects of habitat conversion on activity patterns of phyllostomid bats in lowland Amazonian rainforest. *Journal of Mammalogy* 90:210-221.

Presley SJ, Willig MR, Saldanha LN, Wunderle JM, Castro-Arellano I. 2009b. Reduced-impact logging has little effect on temporal activity of frugivorous bats (Chiroptera) in lowland Amazonia. *Biotropica* 41:369-378.



RECENT PUBLICATIONS

- Adler, F. R., C. A. Clay, and E. M. Lehmer. 2008. The role of heterogeneity in the persistence and prevalence of Sin Nombre virus in deer mice. *Am. Nat.*, 172:855-867.
- Alvarez-Castaneda, S. T. and N. Gonzalez-Ruiz. 2009. *Peromyscus levipes* (Rodentia: Cricetidae). *Mammal. Sp.*, 824:1-6.
- Armien, A. G., B. Armien, F. Koster, J. M. Pascale, M. Avila, P. Gonzalez, M. de la Cruz, Y. Zaldivar, Y. Mendoza, F. Gracia, B. Hjelle, S.-J. Lee, T. L. Yates, and J. Salazar-Bravo. 2009. Hantavirus infection and habitat associations among rodent populations in agroecosystems of Panama: implications for human disease risk. *Am. J. Trop. Med. Hyg.*, 81:59-66.
- Bartel, R. A., F. F. Knowlton, and L. C. Stoddart. 2008. Long-term patterns in mammalian abundance in northern portions of the Great Basin. *J. Mammal.*, 89:1170-1183.
- Battisti, J. M., J. L. Bono, P. A. Rosa, M. E. Schrupf, T. G. Schwan, and P. F. Policastro. 2008. Outer surface protein A protects Lyme disease spirochetes from acquired host immunity in the tick vector. *Infect. Immunity*, 76:5228-5237.
- Black, W. C. IV, J. B. Doty, M. T. Hughes, B. J. Beaty, and C. H. Calisher. 2009. Temporal and geographic evidence for evolution of Sin Nombre virus using molecular analyses of viral RNA from Colorado, New Mexico and Montana. *Virology*, 6:102
<http://www.virologyj.com/content/6/1/102>
- Booher, C. M and W. R. Hood. 2009. Calcium intake, bone metabolism and reproductive output of white-footed mice (*Peromyscus leucopus*). *Integr. Comp. Biol.*, 49:E202. Meeting abstract.
- Briggs, J. S., S. B. V. Wall, and S. H. Jenkins. 2009. Forest rodents provide directed dispersal of Jeffrey pine seeds. *Ecology*, 90:675-687.
- Brittain, R. A., V. J. Meretsky, J. A. Gwinn, J. G. Hammond, J. K. Riegel. 2009. Northern saw-whet owl (*Aegolius acadicus*) autumn migration magnitude and demographics in south-central Indiana. *J. Raptor Res.*, 43:199-209.
- Broussard, D. R. J. L. Robertson, and T. J. Evans II. 2009. Age-related decline in reproductive sensitivity to inhibition by short photoperiod in *Peromyscus leucopus*. *J. Mammal.*, 90:32-39.

- Busch, J. D., P. M. Waser, and J. A. DeWoody. 2008. Characterization of expressed class II MHC sequences in the banner-tailed kangaroo rat (*Dipodomys spectabilis*) reveals multiple DRB loci. *Immunogenetics*, 60:677-688.
- Campbell, P., R. L. Reep, M. L. Stoll, A. G. Ophir, and S. M. Phelps. 2009. Conservation and diversity of Foxp2 expression in muroid rodents: functional implications. *J. Comp. Neurol.*, 512:84-100.
- Castro-Arellano, I., T. E. Lacher Jr. 2009. Temporal niche segregation in two rodent assemblages of subtropical Mexico. *J. Trop. Ecol.*, 25:593-603.
- Castro-Arellano, I., G. Suzan, R. F. Leon, R. M. Jimenez, and T. E. Lacher Jr. 2009. Survey for antibody to hantaviruses in Tamaulipas, Mexico. *J. Wildl. Dis.*, 45:207-212.
- Chappell, M. A. and E. M. Dlugosz. 2009. Aerobic capacity and running performance across a 1.6 km altitude difference in two sciurid rodents. *J. Exp. Biol.*, 212:610-619.
- Chavez, C. and G. Ceballos. 2009. Implications for conservation of the species diversity and population dynamics of small mammals in an isolated reserve in Mexico City. *Nat. Areas J.*, 29:27-41.
- Clay, C. A., E. M. Lehmer, A. Previtali, S. St. Jeor, and M. D. Dearing. 2009. Contact heterogeneity in deer mice: implications for Sin Nombre virus transmission. *Proc. R. Soc. B-Biol. Sci.*, 276:1305-1312.
- Clay, C. A., E. M. Lehmer, S. St. Jeor, and M. D. Dearing. 2009. Sin Nombre virus and rodent species diversity: a test of the dilution and amplification hypotheses. *PLoS ONE*, 4(7):e6467. doi:10.1371/journal.pone.0006467
- Clay, C. A., E. M. Lehmer, S. St. Jeor, and M. D. Dearing. 2009. Testing mechanisms of the dilution effect: deer mice encounter rates, Sin Nombre virus prevalence and species diversity. *EcoHealth*, DOI: 10.1007/s10393-009-0240-2
- Davis, R. M., E. Cleugh, R. T. Smith, and C. L. Fritz. 2008. Use of a chitin synthesis inhibitor to control fleas on wild rodents important in the maintenance of plague, *Yersinia pestis* in California. *J. Vector Ecol.*, 33:278-284.
- Dearing, M. D., M. A. Previtali, J. D. Jones, P. W. Ely, and B. A. Wood. 2009. Seasonal variation in Sin Nombre virus infections in deer mice: preliminary results. *J. Wildl. Dis.*, 45:430-436.
- de Jong, T. R., M. Chauke, B. N. Harris, and W. Saltzman. 2009. From here to paternity: neural correlates of the onset of paternal behavior in California mice (*Peromyscus californicus*). *Horm. Behav.*, 56:220-231.

- de Sousa, R. L. M., M. L. Moreli, A. A. Borges, G. M. Campos, M. C. Livonesi, L. T. M. Figueriredo, and A. A. Pinto. 2008. Natural host relationships and genetic diversity of rodent-associated hantaviruses in southeastern Brazil. *Intervirology*, 51:299-310.
- Dizney, L. J. and L. A. Ruedas. 2009. Increased host species diversity and decreased prevalence of Sin Nombre virus. *Emerg. Infect. Dis.*, 15:1012-1018.
- Easterbrook, J. D. and S. L. Klein. 2008. Immunological mechanisms mediating hantavirus persistence in rodent reservoirs. *PLoS*, 4:e1000172
www.plospathogens.org
- Eisen, R. J., J. L. Holmes, A. M. Schotthoefer, S. M. Vetter, J. A. Montenieri, and K. L. Gage. 2008. Demonstration of early-phase transmission of *Yersinia pestis* by the mouse flea *Aetheca wagneri* (Siphonaptera: Ceratophylidae), and implications for the role of deer mice as enzootic reservoirs. *J. Med. Entomol.*, 45:1160-1164.
- Erickson, H. J., C. L. Aldridge, and N. T. Hobbs. 2009. Progress report: Stratton Ecological Research Site – an experimental approach to assess effects of various grazing treatments on vegetation and wildlife communities across managed burns and habitat controls. US Geol. Survey Report 2009-1016:i-iii, 1-15.
- Fuxiager, M. J., G. Mast, E. A. Becker, and C. A. Marler. 2009. The ‘home advantage’ is necessary for a full winner effect and changes in post-encounter testosterone. *Horm. Behav.*, 56:214-219.
- Gatewood, A. G., K. A. Liebman, G. Vourc’h, J. Bunikis, S. A. Hamer, R. Cortinas, F. Melton, P. Cislo, U. Kitron, J. Tsao, A. G. Barbour, D. Fish, and M. A. Diuk-Wasser. 2009. Climate and tick seasonality are predictors of *Borrelia burgdorferi* genotype distribution. *Appl. Environ. Microbiol.*, 75:2476-2483.
- Geluso, K. 2009. Distributional records for seven species of mammals in southern New Mexico. *Occasional Papers, Museum Texas Tech Univ.*, 287:1-7.
- Gering, E. J., J. C. Opazo, and J. F. Storz. 2009. Molecular evolution of cytochrome b in high- and low-altitude deer mice (genus *Peromyscus*). *Heredity*, 102:226-235.
- Gleason, E. D., M. J. Fuxjager, T. O. Oyegbile, and C. A. Marler. 2009. Testosterone release and social context: when it occurs and why. *Front Neuroendocrinol.*, 30:460-469.
- Grear, D. A., S. E. Perkins, and P. J. Hudson. 2009. Does elevated testosterone result in increased exposure and transmission of parasites? *Ecol. Lett.*, 12:528-537.
- Heideman, P. D. and J. T. Pittman. 2009. Evolution of neuroendocrine mechanisms that regulate reproduction in white-footed mice (*Peromyscus leucopus*). *Integr. Comp. Biol.*, 49:E73. Meeting abstract.

- Hopton, M. E., G. N. Cameron, M. J. Cramer, M. Polak, and G. W. Uetz. 2009. Live animal radiography to measure developmental instability in populations of small mammals after a natural disaster. *Ecol. Indicators*, 9:883-891.
- Hortelano-Moncada, Y., F. A. Cervantes, and A. Trejo-Ortiz. 2009. Wild mammals of Reserva Ecologica del Pedregal de San Angel, Ciudad Universitaria, UNAM, Mexico, DF. *Rev. Mex. Biodivers.*, 80:507-520.
- Huggard, D. J. and A. Arsenault. 2009. Conifer seed predation in harvested and burned dry Douglas-fir forests in southern British Columbia. *Can. J. For. Res.-Rev. Can. Rech. For.*, 39:1548-1556.
- Ivanova, L., I. Christova, V. Neves, M. Aroso, L. Meirelles, D. Brisson, and M. Gomes-Solecki. 2009. Comprehensive seroprofiling of sixteen *B. burgdorferi* OspC: implications for Lyme disease diagnostics design. *Clin. Immunol.*, 132:393-400.
- Johnson, E. M., R. J. Panciera, K. E. Allen, M. E. Sheets, J. D. Beal, S. A. Ewing, and S. E. Little. 2009. Alternate pathway of infection with *Hepatozoon americanum* and the epidemiologic importance of predation. *J. Vet. Intern. Med.*, Aug 26 [Epub ahead of print]
- Johnson, K. E., L. D. Knopper, D. C. Schneider, C. A. Ollson, and K. J. Reimer. 2009. Effects of local point source polychlorinated biphenyl (PCB) contamination on bone mineral density in deer mice (*Peromyscus maniculatus*). *Sci. Total Environ.*, 407:5050-5055.
- Johnston, A. N. and R. G. Anthony. 2008. Small-mammal microhabitat associations and response to grazing in Oregon. *J. Wildlife Manage.*, 72:1736-1746.
- Kalcounis-Rueppell, M. C. and T. R. Spoon. 2009. *Peromyscus boylii* (Rodentia: Cricetidae). *Mammal. Sp.*, 838:1-14.
- Keesing, F., J. Brunner, S. Duerr, M. Killilea, K. LoGiudice, K. Schmidt, H. Vuong, and R. S. Ostfeld. 2009. Hosts as ecological traps for the vector of Lyme disease. *Proc. R. Soc. B.*, 276:3911-3919.
- Kingsley, E. P., M. Manceau, C. D. Wiley, H. E. Hoekstra. 2009. Melanism in *Peromyscus* is caused by independent mutation in *Agouti*. *PLoS ONE*, 4(7):e6435. doi:10.1371/journal.pone.0006435
- Klenner, W. and T. P. Sullivan. 2009. Partial and clearcut harvesting of dry Douglas-fir forests: implications for small mammal communities. *For. Ecol. Manage.*, 257:1078-1086.

- Klinkenborg, V. 2009. Efforts at protection are celebrated-and scorned. Essay In: National Geographic, Jan.:82-107.
- Korff, S., D. J. Stein, and B. H. Harvey. 2009. Cortico-striatal cyclic AMP-phosphodiesterase-4 signalling and stereotypy in the deer mouse: attenuation after chronic fluoxetine treatment. *Pharmacol. Biochem. Behav.*, 92:514-520.
- Korva, M., D. Duh, A. Saksida, T. Trilar, and T. Avsic-Zupanc. 2009. The hantaviral load in tissues of naturally infected rodents. *Microbes Infect.*, 11:344-351.
- Labinskyy, N., P. Mukhopadhyay, J. Toth, G. Szalai, M. Veres, G. Losonczy, J. T. Pinto, P. Pacher, P. Ballabh, A. Podlutsky, S. N. Austad, A. Csiszar, and Z. Ungvari. 2009. Longevity is associated with increased vascular resistance to high glucose-induced oxidative stress and inflammatory gene expression in *Peromyscus leucopus*. *Am. J. Physiol. Heart Circ. Physiol.*, 296:946-956.
- Lambert, K. 2009. Rodent world holds clues about ours. Richmond Times-Dispatch By Karin Kapsidelis Published: February 14, 2009
http://www2.timesdispatch.com/rtd/news/local/education/article/MICE14_20090213-222728/206471/
http://static.mgnetwork.com/rtd/slideshows/20090214_MICE/index.html
- Lazzari, V., C. Charles, P. Tafforeau, M. Vianey-Liaud, J-P. Aguilar, J-J. Jaeger, J. Michaux, and L. Viriot. 2008. Mosaic convergence of rodent detitions. *PLoS One*, 3:e3607. www.plosone.org
- Lemaitre, J., D. Fortin, P. O. Montiglio, and M. Darveau. 2009. Bot fly parasitism of the red-backed vole: host survival, infection risk, and population growth. *Oecologia*, 159:283-294.
- Linnen, C. R., E. P. Kingsley, J. D. Jensen, and H. E. Hoekstra. 2009. On the origin and spread of an adaptive allele in deer mice. *Science*, 325:1095-1098.
- Lobo, N., M. Duong, and J. S. Millar. 2009. Conifer-seed preferences of small mammals. *Can. J. Zool.*, 87:773-780.
- LoGiudice, K., S. T. K. Duerr, M. J. Newhouse, K. A. Schmidt, M. E. Killilea, and R. S. Ostfeld. 2008. Impact of host community composition on Lyme disease risk. *Ecology*, 89:2841-2849.
- Louzada, S., A. Paco, S. Kubickova, F. Adegas, H. Guedes-Pinto, J. Rubes, and R. Chaves. 2008. Different evolutionary trails in the related genomes *Cricetus cricetus* and *Peromyscus eremicus* (Rodentia, Cricetidae) uncovered by orthologous satellite DNA repositioning. *Micron*, 39:1149-1155.

- Louzada, S., A. Vieira-da-Silva, F. Adegá, H. Guedes-Pinto, and R. Chaves. 2009. Mouse and rat genomes uncover the chromosome restructuring in *Peromyscus eremicus* (Cricetidae, Rodentia)-Chromosomes 1 and 5 in focus. *Chromosome Res.*, 17:168-169. Meeting Abstract
- Luis, J., L. Ramirez, A. Carmona, G. Ortiz, J. Delgado, and R. Cardenas. 2009. Paternal behavior and testosterone plasma levels in the Volcano mouse *Neotomodon alstoni* (Rodentia: Muridae). *Rev. Biol. Trop.*, 57:433-439.
- Luong, L. T., D. A. Grear, and P. J. Hudson. 2009. Male hosts are responsible for the transmission of a trophically transmitted parasite, *Pterygodermatites peromysci*, to the intermediate host in the absence of sex-biased infection. *Int. J. Parasitol.*, 39:1263-1268.
- Mabry, K. E. and J. A. Stamps. 2008. Searching for a new home: decision making by dispersing brush mice. *Am. Nat.*, 172:625-634.
- McCaffrey, R. E., M. C. Wallace, and J. D. Ray. 2009. Small mammals and ground-dwelling invertebrates associated with active and controlled colonies of black-tailed prairie dogs (*Cynomys ludovicianus*). *Southwest. Naturalist*, 54:300-306.
- McEachern, M. B., R. L. McElreath, D. H. Van Vuren, and J. M. Eadie. 2009. Another genetically promiscuous 'polygynous' mammal: mating system variation in *Neotoma fuscipes*. *Anim. Behav.*, 77:449-455.
- McPhee, M. E., A. E. Ribbeck, and R. E. Johnston. 2009. Male golden hamsters (*Mesocricetus auratus*) are more reactive than females to a visual predator cue. *J. Ethol.*, 27:137-141.
- Miller, J. R. and M. D. Engstrom. 2008. The relationships of major lineages within Peromyscine rodents: a molecular phylogenetic hypothesis and systematic reappraisal. *J. Mammal.*, 89:1279-1295.
- Mlynarski, E. E., C. J. Obergfell, G. J. Szalai, and R. J. O'Neill. 2009. Epigenetic basis for cancer susceptibility in a *Peromyscus* model. *Chromosome Res.*, 17:560. Meeting abstract.
- Mullen, L. M., S. N. Vignieri, J. A. Gore, and H. E. Hoekstra. 2009. Adaptive basis of geographic variation genetic, phenotypic and environmental differences among beach mouse populations. *Proc. R. Soc. B.*, 276:3809-3818.
- Nietro, N. C. and J. E. Foley. 2009. Meta-analysis of coinfection and coexposure with *Borrelia burgdorferi* and *Anaplasma phagocytophilum* in humans, domestic animals, wildlife, and *Ixodes ricinus*-complex ticks. *Vector-Borne Zoonotic Dis.*, 9:93-101.

- Ollson, C. A., I. Koch, P. Smith, L. D. Knopper, C. Hough, and K. J. Reimer. 2009. Addressing arsenic bioaccessibility in ecological risk assessment: a novel approach to avoid overestimating risk. *Environ. Toxicol. Chem.*, 28:668-675.
- Oriel, R. C., C. D. Wiley, M. J. Dewey, and P. B. Vrana. 2008. Adaptive genetic variation, stress and glucose regulation. *Dis. Mod. Mech.*, 1:255–263.
- Orrock, J. L. and B. F. Allan. 2008. Sin Nombre virus infection in deer mice, Channel Islands, California. *Emerg. Infect. Dis.*, 14:1965-1966.
- Orrock, J. L. and B. J. Danielson. 2009. Temperature and cloud cover, but not predator urine, affect winter foraging of mice. *Ethology*, 115:641-648.
- Paco, A., F. Adegá, H. Guedes-Pinto, and R. Chaves. 2008. Molecular analysis and physical distribution of LINE-1 sequences in three rodent species, *Cricetus cricetus*, *Peromyscus eremicus* (Cricetidae family) and *Praomys tullbergi* (Muridae family). *Chromosome Res.*, 16:1047. Meeting abstract.
- Paco, A., F. Adegá, H. Guedes-Pinto, and R. Chaves. 2009. Hidden heterochromatin: characterization in the rodentia species *Cricetus cricetus*, *Peromyscus eremicus* (Cricetidae) and *Praomys tullbergi* (Muridae). *Genetics Mol. Biol.*, 32:58-68.
- Paniagua, L. L. and J. J. Morrone. 2009. Do the Oaxacan Highlands represent a natural biotic unit? A cladistic biogeographical test based on vertebrate taxa. *J. Biogeogr.*, 36:1939-1944.
- Patz, J. A., S. H. Olson, C. K. Uejio, and H. K. Gibbs. 2008. Disease emergence from global climate and land use change. *Med. Clin. N. Amer.*, 92:1473-1491.
- Pennisi, E. 2009. How beach life favors blond mice. *News Focus Science*, 325:1330-1333.
- Pries, A. J., L. C. Branch, and D. L. Miller. 2009. Impact of hurricanes on habitat occupancy and spatial distribution of beach mice. *J. Mammal.*, 90:841-850.
- Ramm, S. A., L. McDonald, J. L. Hurst, R. J. Beynon, and P. Stockley. 2009. Comparative proteomics reveals evidence for evolutionary diversification of rodent seminal fluid and its functional significance in sperm competition. *Mol. Biol. Evol.*, 26:189-198.
- Randa, L. A., D. M. Cooper, P. L. Meserve, and J. A. Yunker. 2009. Prey switching of sympatric canids in response to variable prey abundance. *J. Mammal.*, 90:594-603.

- Ransome, D. B., P. M. F. Lindgren, M. J. Waterhouse, H. M. Armleder, and T. P. Sullivan. 2009. Small-mammal response to group-selection silvicultural systems in Engelmann spruce – subalpine fir forests 14 years postharvest. *Can. J. Forest Res.*, 39:1698-1708.
- Rayfield, B., A. Moilanen, and M. J. Fortin. 2009. Incorporating consumer-resource spatial interactions in reserve design. *Ecol. Model.*, 220:725-733.
- Rezende, E. L., K. A. Hammond, and M. A. Chappell. 2009. Cold acclimation in *Peromyscus*: individual variation and sex effects in maximum and daily metabolism, organ mass and body composition. *J. Exp. Biol.*, 212:2795-2802.
- Richens, T., A. D~N Palmer, J. Prescott, and T. Shountz. Genomic organization and phylogenetic utility of deer mouse (*Peromyscus maniculatus*) lymphotoxin-alpha and lymphotoxin-beta. *BMC Immunol.*, 9:62. www.biomedcentral.com/1471-2172/9/62
- Ridley, M. 2009. Modern Darwins. *National Geographic*, February 2009.
- Rodas, L. R., C. A. Jennison, D. B. Hall, G. W. Barrett. 2009. Luring small mammals: a levels-of-organization perspective. *Southeast. Nat.*, 8:387-398.
- Rogers, E. A., S. V. Abdunnur, J. V. McDowell, and R. T. Marconi. 2009. Comparative analysis of the properties and ligand binding characteristics of CspZ, a factor H binding protein, derived from *Borrelia burgdorferi* isolates of human origin. *Infect. Immunity*, 77:4396-4405.
- Ronconi, R. A. and J. M. Hipfner. 2009. Egg neglect under risk of predation in Cassin's auklet (*Ptychoramphus aleuticus*). *Can. J. Zool.*, 87:415-421.
- Rudenko, N., M. Golovchenko, L. Grubhoffer, and J. H. Oliver. 2009. *Borrelia carolinensis* sp nov., a new (14th) member of the *Borrelia burgdorferi* sensu lato Complex from the southeastern region of the United States. *J. Clin. Microbiol.*, 47:134-141.
- Sanchez-Hernandez, C., G. D. Schnell, M. de Lourdes Romero-Almaraz. 2009. *Peromyscus perfulvus* (Rodentia: Cricetidae). *Mammal. Sp.*, 833:1-8.
- Santolo, G. M. 2009. Small mammals collected from a site with elevated selenium concentrations and three reference sites. *Arch. Environ. Contam. Toxicol.*, Apr 17. [Epub ahead of print]
- Satoh, K. and F. Iwaku. 2009. Structure and direction of jaw adductor muscles as herbivorous adaptations in *Neotoma mexicana* (Muridae, Rodentia). *Zoomorphology*, 128:339-348.

- Schauber, E. M., M. J. Connors, B. J. Goodwin, C. G. Jones, and R. S. Ostfeld. 2009. Quantifying a dynamic risk landscape: heterogeneous predator activity and implications for prey persistence. *Ecology*, 90:240-251.
- Schwanz, L. E., D. Brisson, M. Gomes-Solecki, and R. S. Ostfeld. 2009. The impact of the spirochete *Borrelia burgdorferi* on white-footed mice: implications for the ecology of Lyme disease. *Integr. Comp. Biol.*, 49:E153. Meeting abstract.
- Sears, M. W., J. P. Hayes, M. R. Banta, and D. McCormic. 2009. Out in the cold: physiological capacity influences behaviour in deer mice. *Funct. Ecol.*, 23:774-783.
- Shahid, A., D. E. Garneau, and T. S. McCay. 2009. Selection of seeds of common native and non-native plants by granivorous rodents in the northeastern United States. *Am. Midl. Nat.*, 162:207-212.
- Shi, Y., R. Buffenstein, S. N. Austad, and H. VanRemmen. 2008. Skin-derived fibroblasts of house mouse (*Mus musculus*) and white-footed mouse (*Peromyscus leucopus*) show distinct metabolic profiles in culture. *Free Rad. Biol. Med.*, 45:S100. Meeting abstract.
- Slowick, T. J., and R. S. Lane. 2009. Feeding preferences of the immature stages of three western North American Ixodid ticks (*Acair*) for avian, reptilian, or rodent hosts. *J. Med. Entomol.*, 46:115-122.
- Smith, J. N., M. A. Espino, J. Liu, N. A. Romero, S. B. Cox, and G. P. Cobb. 2009. Multigenerational effects in deer mice (*Peromyscus maniculatus*) exposed to hexahydro-1,3,5-trinitroso-1,3,5-triazine (TNX). *Chemosphere*, 75:910-914.
- Stacey, C. A. and D. M. Post. 2009. Effects of disturbance by humans on small mammals in a Chihuahuan Desert ecosystem. *Southwest. Naturalist*, 54:272-278.
- Steiner, C. C., H. Rompler, L. M. Boettger, T. Schoneberg, and H. E. Hoekstra. 2009. The genetic basis of phenotypic convergence in beach mice: similar pigment patterns but different genes. *Mol. Biol. Evol.*, 26:35-45.
- Storz, J. F., A. M. Runck, S. J. Sabatino, J. K. Kelly, N. Ferrand, H. Morriyama, R. E. Weber, and A. Fago. 2009. Evolutionary and functional insights into the mechanism underlying high-altitude adaptation of deer mouse hemoglobin. *PNAS*, 106:14450-14455.
- Suarez-Gracida, C. G. and S. T. Alvarez-Castaneda. 2009. Physical and biological variables related to habitat preferences of rodents. *Biodivers. Conserv.*, 18:2779-2797.
- Sullivan, T. P. and D. S. Sullivan. 2009. Are linear habitats in agrarian landscapes source areas of beneficial or pest rodents? *Agric. Ecosyst. Environ.*, 129:52-56.

- Tanimura, Y., F. C. Ogoegbunam, M. H. Lewis. 2009. Amphetamine-induced sensitization and spontaneous stereotypy in deer mice. *Pharmacol., Biochem. Behav.*, 92:670-675.
- Tannenbaum, L. V., B. H. Thran, and K. J. Williams. 2009. Testing the limits of rodent sperm analysis: azoospermia in an otherwise healthy wild rodent population. *Arch. Environ. Contam. Toxicol.*, 56:157-164.
- Taraborelli, P., N. Borruel, and A. Mangeaud. 2009. Ability of murid rodents to find buried seeds in the Monte Desert. *Ethology*, 115:201-209.
- Timonin, M. E., B. S. Cushing, and K. E. Wynne-Edwards. 2008. In three brain regions central to maternal behaviour, neither male nor female *Phodopus* dwarf hamsters show changes in oestrogen receptor alpha distribution with mating or parenthood. *Neuroendocrinol.*, 20:1301-1309.
- Trainor, B. C., C. L. Sisk, and R. J. Nelson. 2009. Hormones and the development and expression of aggressive behavior. In: D. W. Pfaff, A. P. Arnold, A. M. Etgen, S. E. Fahrbach and R. T. Rubin, editors. *Hormones, Brain and Behavior*, 2nd edition, Vol 1. San Diego: Academic Press:167-203.
- Ungvari, Z., B. F. Krasnikov, A. Csiszar, N. Labinskyy, P. Mukhopadhyay, P. Pacher, A. J. L. Cooper, N. Podlutskaya, S. N. Austad, and A. Podlutsky. 2008. Testing hypotheses of aging in long-lived mice of the genus *Peromyscus*: association between longevity and mitochondrial stress resistance, *ROS* detoxification pathways, and DNA repair efficiency. *AGE*, 30:121-133.
- Vandegrift, K. J. and P. J. Hudson. 2009. Could parasites destabilize mouse populations? The potential role of *Pterygodermatites peromysci* in the population dynamics of free-living mice, *Peromyscus leucopus*. *Int. J. Parasitol.*, 39:1253-1262.
- Vandegrift, K. J. and P. J. Hudson. 2009. Response to enrichment, type and timing: small mammals vary in their response to a springtime cicada but not a carbohydrate pulse. *J. Anim. Ecol.*, 78:202-209.
- Van Wynsberghe, N. R., S. B. Canto-Lara, E. I. Sosa-Bibiano, N. A. Rivero-Cardenas, and F. J. Andrade-Narvaez. 2009. Comparison of small mammal prevalence of *Leishmania (Leishmania) mexicana* in five foci of Cutaneous Leishmaniasis in the state of Campeche, Mexico. *Rev. Inst. Med. Trop. S. Paulo*, 51:87-94.
- Wang, G. M., J. Wolff, S. Vessey, N. Slade, J. Witham, J. Merritt, M. Hunter, and S. Elias. 2009. Comparative population dynamics of *Peromyscus* in North America: influences of climate, food, and density dependence. *Popul. Ecol.*, 51:133-142.
- Weber, J. N. and H. E. Hoekstra. 2009. The evolution of burrowing behaviour in deer mice (genus *Peromyscus*). *Anim. Behav.*, 77:603-609.

- Weber, J. N., M. B. Peters, O. V. Tsyusko, C. R. Linnen, C. Hagen, N. A. Schable, T. D. Tuberville, A. M. McKee, S. L. Lance, K. L. Jones, H. S. Fisher, M. J. Dewey, H. E. Hoekstra, and T. C. Glenn. 2009. Five hundred microsatellite loci for *Peromyscus*. *Conserv. Genet.*, In Press.
- Wiley, C. D., H. H. Matundan, A. R. Duselis, A. T. Isaacs, and P. B. Vrana. 2008. Patterns of hybrid loss of imprinting reveal tissue- and cluster-specific regulation. *PLoS One*, 3:e3572. www.plosone.org
- Williams, S. C., J. S. Ward, T. E. Worthley, and K. C. Stafford III. 2009. Managing Japanese barberry (Ranunculales: Berberidaceae) infestations reduces blacklegged tick (Acari: Ixodidae) abundance and infection prevalence with *Borrelia burgdorferi* (Spirochaetales: Spirochaetaceae). *Environ. Entomol.*, 38:977-984.
- Williams, S. H., E. Peiffer, and S. Ford. 2009. Gape and bite force in the rodents *Onychomys leucogaster* and *Peromyscus maniculatus*: does jaw-muscle anatomy predict performance? *J. Morphology*, [Epub ahead of print]
- Wilson, A. J., U. Gelin, M. C. Perron, and D. Reale. 2009. Indirect genetic effects and the evolution of aggression in a vertebrate system. *Proc. Boil. Sci.*, 276:533-541.
- Wilson, J. A., J. L. Parsons, and E. E. Jorgensen. 2009. Protein nutrition of southern plains small mammals: immune response to variation in maternal and offspring dietary nitrogen. *Physiol. Biochem. Zool.*, 82:170-180.
- Workman, J. L., S. L. Bowers, and R. J. Nelson. 2009. Enrichment and photoperiod interact to affect spatial learning and hippocampal dendritic morphology in white-footed mice (*Peromyscus leucopus*). *Eur. J. Neurosci.*, 29:161-170.
- Wu, H. M., C. C. White, J.P. Isanhart, T. J. McBride, T. J. Kavanagh, and M. J. Hooper. 2009. Optimization and application of glutamate cysteine ligase measurement in wildlife species. *Ecotoxicol. Environ. Safety*, 72:572-578.
- Wu, H., T. J. McBride, J. P. Isanhart, S. B. Cox, and M. J. Hooper. 2009. Responses of glutamate cysteine ligase and glutathione to oxidants in deer mice (*Peromyscus maniculatus*). *Ecotoxicol. Environ. Safety*, 72:1572-1578.
- Zelditch, M. L., A. R. Wood, R. M. Bonett, and D. L. Swiderski. 2009. Modularity of the rodent mandible: integrating bones, muscles, and teeth. *Evol. Dev.*, 10:756-768.
- Zelditch, M. L., A. R. Wood, and D. L. Swiderski. 2009. Building developmental integration into functional systems: function-induced integration of mandibular shape. *Evol. Biol.*, 36:71-87.

Ziegler, P. E., S. E. Wade, J. Oliver, S. L. Schaaf, and H. O. Mohammed. 2009. Factors associated with the risk of giardia infection in two sympatric mouse species of the genus *Peromyscus*. *J. Parasitol.*, April 14:1 [Epub ahead of print]

Zwolak, R. 2009. A meta-analysis of the effects of wildfire, clearcutting, and partial harvest on the abundance of North American small mammals. *For. Ecol. Management*, 258:539-545.

* * *